

ANALYTICAL STUDY OF LOW TEMPERATURE PHASE OF 3D LGT IN THE PLAQUETTE FORMULATION

OLEG BORISENKO, SERGEI VOLOSHIN

*Institute for Theoretical Physics, Ukrainian Academy of
Sciences, 03143 Kiev, Ukraine*

AND

MANFRIED FABER

*Atominstitut der österreichischen Universitäten,
Arbeitsgruppe Kernphysik, TU Wien, A-1040 Vienna, Austria*

Abstract. We develop an analytical approach for non-abelian gauge models within the plaquette representation where the plaquette matrices play the role of the fundamental degrees of freedom. We start from the original Batrouni formulation and show how it can be modified in such a way that each non-abelian Bianchi identity contains two connectors instead of four. Using this representation we construct the low-temperature expansion for $U(1)$ and $SU(N)$ models on a finite lattice and discuss its uniformity in the volume. Next, we derive a dual representation for the 't Hooft loop in the $SU(2)$ model and describe monopoles in the maximal axial gauge.

1. Plaquette formulation

Lattice gauge theory (LGT) can be formulated in many equivalent ways. The original Wilson formulation is given in terms of group valued matrices on links as fundamental degrees of freedom [1]. The partition function reads

$$Z = \int DU \exp\{-\beta S[U_\mu(x)]\} , \quad (1)$$

where S is the standard Wilson action and the integral is calculated over the Haar measure on the group at every link of the lattice. Several years later

Presented by O. Borisenko at the NATO Advanced Research Workshop “Confinement, Topology, and other Non-Perturbative Aspects of QCD”, January 21–27, 2002, Stará Lesná, Slovakia.

the dual representation for abelian LGT's was constructed in [2]. Extensions to nonabelian groups have been proposed in nineties in [3]-[4]. A closely related approach is so-called plaquette representation invented in the continuum theory by M. Halpern [5] and extended to the lattice models in [6]. In this representation the plaquette matrices play the role of the dynamical degrees of freedom and satisfy certain constraints expressed through Bianchi identities. Both dual and plaquette formulations are not so popular in the case of nonabelian models, probably due to their complexity. In the nonabelian case above mentioned constraints appear to be highly-nonlocal and this fact makes an analytical study of the model rather difficult. In spite of this complexity the plaquette formulation has certain advantages over the standard one [6]: duality transformations, Coulomb-gas representation, strong coupling expansion and mean-field approximation look more natural and simpler in the plaquette formulation. Nevertheless we believe the main advantage of this formulation lies in its applications to the low-temperature region. Let V_p be a plaquette matrix in $SU(N)$ LGT. The rigorous result of [7] asserts that the probability $p(\xi)$ that $\text{Tr}(I - V_p) \geq \xi$ is bounded by

$$p(\xi) \leq \mathcal{O}(e^{-b\beta\xi}) , \quad \beta \rightarrow \infty , \quad b = \text{const} \quad (2)$$

uniformly in the volume. Thus, all configurations with $\xi \geq O(\beta^{-1})$ are exponentially suppressed. The Gibbs measure at large β is strongly concentrated around configurations on which $V_p \approx I$. This property justifies expansion of plaquette matrices around unity when β is sufficiently large.

This is our first motivation to construct a low-temperature expansion of gauge models using the plaquette formulation. The well-known problem of the standard perturbation theory (PT) can be shortly formulated as follows. When the volume is fixed, and in the complete axial gauge the link matrices perform small fluctuations around the unit matrix and the PT works very well producing the asymptotic expansion in inverse powers of β . However, in the thermodynamic limit (TL) system deviates arbitrarily far from the ordered state, no saddle point exists anymore and configurations of link matrices are distributed uniformly in the group space. That there are problems with the conventional PT was proven in [8], where it was shown the PT results depend on the boundary conditions (BC) used to reach the TL. All this rises the question whether PT gives asymptotic expansion uniformly valid in the volume. Fortunately, even in the TL the plaquette matrices are close to unity, the inequality (2) holds and thus provides a basis for the construction of the low-temperature expansion.

Our next motivation is to derive the dual representation of nonabelian models which could be used for their analytical study. Here we formulate such representation for $SU(2)$ LGT and compute dual form of the 't Hooft loop. We also describe monopoles of the maximal axial gauge.

Now we outline the derivation of our plaquette formulation (all technical details will be given in a forthcoming publication [9]). Consider a 3D lattice with free BC on the link gauge matrices. We make a change of variables in the partition function (1) $V_p = U_\mu(x)U_\nu(x + \mu)U_\mu^\dagger(x + \nu)U_\nu^\dagger(x)$. The partition function gets the form

$$Z = \int \prod_p dV_p \exp \left[\beta \sum_p \text{Re Tr} V_p \right] J(V_p) , \quad (3)$$

$$J(V_p) = \int \prod_l dU_l \prod_p \delta \left(V_p^\dagger \prod_{l \in p} U_l - I \right) = \prod_c J(V_c) . \quad (4)$$

The last equations are rather formal, for we have to specify the order of multiplication of nonabelian matrices. To do that we define in a cube a vertex A and separated by the space diagonal a vertex B , see Fig.1. This assignment we extend for all neighbouring cubes and finally for the whole lattice. The plaquette matrices V_p we insert at the vertices A or B of each plaquette attached to A or B , correspondingly. Next, take a path connecting vertices A and B by three links, as shown in Fig.1. Then the matrix V_c entering the Bianchi identity for the cube c is of the form

$$V_c = \exp \left[i\sigma^k \omega_k(c) \right] = \left(\prod_{p \in A} V_p \right) C \left(\prod_{p \in B} V_p \right) C^\dagger , \quad C = U_1 U_2 U_3^\dagger . \quad (5)$$

$\prod_{p \in (A,B)}$ means an appropriately ordered product over three plaquettes of the cube attached to the vertex A or B . The matrix C defines a parallel transport of vertex B into vertex A and is called connector. The connector C plays a crucial role in the nonabelian Bianchi identity. Its path can, in principle be chosen in an arbitrary way but the possible choices of $\prod_{p \in A}$, $\prod_{p \in B}$ and C must fit together. We choose the structure of connectors as shown in Fig.1. As is seen from Fig.1 there are four types of elementary




Figure 1. Structure of connectors on the lattice.

cubes with different types of connector pathes. The next steps in the derivation are as follows:

I). Map one of the space-like plaquette delta's to a delta-function for the cube $J(V_c)$ with V_c given in (5).

II). Choose the maximal axial gauge in the form

$$U_3(x, y, z) = U_2(x, y, 0) = U_1(x, 0, 0) = I . \quad (6)$$

To simplify connectors one can fix the Dirichlet BC in the plane $z = 0$.

III). Integrate out all link variables following [6].

In what follows we pass to the dual lattice where cubes become sites and so on. The partition function takes the form

$$Z = \int \prod_l dV_l \exp[\beta \sum_l \text{Re Tr} V_l] \prod_{i=1}^4 \prod_{x(i)} J(V_x^{(i)}) , \quad (7)$$

where $\prod_{x(i)}$ runs over all sites with i -th type of connector pathes and $J(V_x^{(i)})$ is an $SU(N)$ delta-function given by

$$J(V_x) = \sum_r d_r \chi_r(V_x) , \quad (8)$$

where the sum over r is a sum over all representations of $SU(N)$, χ_r is character of r -th representation and $d_r = \chi_r(I)$.

To write down exact expressions we choose coordinate system on the dual lattice as in Fig.1. Then, e.g. the connector for the cube $B_1 A$ is

$$V_x^{(1)} = V_{l_5}^\dagger V_{l_1} V_{l_6}^\dagger C_{\vec{x}(1)} V_{l_2} V_{l_3} V_{l_4}^\dagger C_{\vec{x}(1)}^\dagger , \quad (9)$$

$$C_{\vec{x}(1)} = \prod_{k=z_i-1}^1 V_{n_2}(x_i, y_i - 1, k) \prod_{p=1}^{z_i-1} V_{n_1}(x_i - 1, y_i, p) . \quad (10)$$

2. Low-temperature expansion

In this section we construct the low-temperature expansion of gauge models in the plaquette formulation. In doing this we follow the strategy developed for the two-dimensional principal chiral models in [10]. The weak-coupling expansion in both models bears many common features. In the abelian case they turn out to be practically identical, while the only, but essential difference in the nonabelian case is the appearance of connectors in gauge models. The lattice PT in temporal gauge ($U_0 = I$) was constructed for abelian and nonabelian models in the standard Wilson formulation by F. Müller and

W. Rühl [11]. In this gauge the PT for nonabelian models displays serious infrared divergences in separate terms of the expansion starting from the order $\mathcal{O}(\beta^{-2})$ for Wilson loops and from $\mathcal{O}(\beta^{-1})$ for the free energy. Although it is expected that all divergences are cancelled in gauge invariant quantities, we are not aware of any proof of the infrared finiteness at higher orders of the expansion. Here we outline the low-temperature expansion for $SU(2)$ LGT (all details and extension to other groups will be given in [9]). We want to expand the partition and correlation functions into asymptotic series whose coefficients are calculated over certain Gaussian measure. Let us consider the standard parameterization for the $SU(2)$ link matrix $V_l = \exp[i\sigma^k \omega_k(l)]$, where $\sigma^k, k = 1, 2, 3$ are Pauli matrices and introduce

$$W_l = \left[\sum_k \omega_k^2(l) \right]^{1/2}, \quad W_x = \left[\sum_k \omega_k^2(x) \right]^{1/2}, \quad (11)$$

where $\omega_k(x)$ is a site angle defined in (5) and has the standard expansion $\omega_k(x) = \omega_k^{(0)}(x) + \omega_k^{(1)}(x) + \omega_k^{(2)}(x) + \dots$ in powers of link angles. The first coefficient can be written down as

$$\omega_k^{(0)}(x) = \omega_k(l_4) + \omega_k(l_5) + \omega_k(l_6) - \omega_k(l_1) - \omega_k(l_2) - \omega_k(l_3). \quad (12)$$

$\omega_k^{(i)}(x)$ for $i \geq 1$ can be computed by making repeated use of the Campbell-Baker-Hausdorff formula. Then, the partition function (7) can be exactly rewritten to the following form [10]

$$Z_{SU(2)} = \int \prod_l \left[\frac{\sin^2 W_l}{W_l^2} \prod_k d\omega_k(l) \right] \exp \left[2\beta \sum_l \cos W_l \right] \prod_x \frac{W_x}{\sin W_x} \\ \prod_x \sum_{m(x)=-\infty}^{\infty} \int \prod_k d\alpha_k(x) \exp \left[-i \sum_k \alpha_k(x) \omega_k(x) + 2\pi i m(x) \alpha(x) \right], \quad (13)$$

where $\alpha(x) = (\sum_k \alpha_k^2(x))^{1/2}$. In derivation of this representation we have used the Poisson formula. In order to perform the weak coupling expansion we make the substitution $\omega_k(l) \rightarrow (2\beta)^{-1/2} \omega_k(l)$, $\alpha_k(x) \rightarrow (2\beta)^{1/2} \alpha_k(x)$ and then expand the integrand of (13) in powers of fluctuations of the link fields. Introduce external sources $h_k(l)$ coupled to the link field $\omega_k(l)$ and $s_k(x)$ coupled to the auxiliary field $\alpha_k(x)$. Then one can get the following expansion

$$Z = \left[1 + \sum_{k=1}^{\infty} \frac{1}{\beta^k} B_k(\partial_h, \partial_s) \right] M(h, s), \quad (14)$$

where B_k are known operators. Up to exponentially small terms the generating functional $M(h, s)$ is given by

$$\exp \left[\frac{1}{4} s_k(x) G_{x,x'} s_k(x') + \frac{i}{2} s_k(x) D_l(x) h_k(l) + \frac{1}{4} h_k(l) G_{ll'} h_k(l') \right], \quad (15)$$

where we introduced link Green functions $G_{ll'}$ and $D_l(x')$ defined as

$$G_{ll'} = 2\delta_{l,l'} - G_{x,x'} - G_{x+n,x'+n'} + G_{x,x'+n'} + G_{x+n,x'} , \quad (16)$$

$$D_l(x') = G_{x,x'} - G_{x+n,x'} . \quad (17)$$

These formulae allow us to calculate the weak coupling expansion of both the free energy and any fixed-distance observable. Extension of this expansion for $SU(N)$ LGT and for the Wilson loop and other observables is straightforward.

Let us give some comments on the property of the expansion and on the infrared problem. One proves that the probability that K links have large fluctuations like $\omega_k(l) \sim \mathcal{O}(\sqrt{\beta})$ is suppressed as $\mathcal{O}(e^{-\text{const} K \pi^2 \beta})$, C is a constant. It shows that large fluctuations are under good control since they are exponentially suppressed with β , and this obviously remains true in the TL. This property is very essential achievement of the expansion in the plaquette representation. No such control exists for the large fluctuations of the link gauge matrices in the TL. It follows from the definition of the link Green function (16) that $|G_{ll'}| \leq 4/3$. Hence, with respect to the Gaussian measure fluctuations of link variables are bounded like

$$|\langle \omega_l \omega_{l'} \rangle_G| = \frac{1}{2} |G_{ll'}| \asymp \frac{\text{const}}{R^3} . \quad (18)$$

This property justifies the low-temperature expansion in powers of fluctuations of link (plaquette) variables, while there is no such justification for the expansion in terms of the original link variables the fluctuations of which are not bounded when $L \rightarrow \infty$. As is seen from the generating functional (15) it depends only on infrared finite Green functions: link Green functions are infrared finite by construction, while the Green function for the free scalar field is finite in any dimension $D \geq 3$.

In spite of all these nice properties, it is far from obvious that the expansion is uniformly valid in the volume. The problem appears in the following way. In [9] we compute the first coefficient of the free energy expansion and prove that it coincides with the standard answer. Nevertheless, that computations show clearly that the source of the troubles are the connectors of the nonabelian Bianchi identities. The low-temperature expansion starts from the abelian Bianchi identity for every cube and goes towards gradual restoration of the full identity with every order of the expansion. Thus, the generating functional contains an abelianized form of the identity without connectors. High-order terms include an expansion of connectors which lead to appearance of infrared divergent terms. Consider, for example the term $\alpha_k(x) \omega_k^1(x) \sim \mathcal{O}(1/\sqrt{\beta})$ in the exponent of (13). $\omega_k^1(x)$ includes the following contributions from connectors $\epsilon_{kmn}(\omega_m(l_1) + \omega_m(l_2) +$

$\omega_m(l_3)) \sum_{l \in C} \omega_n(l)$, where sum over l extends over all links belonging to the connector C . For any finite L one can always find such large β that this sum is of the order $\mathcal{O}(1/\sqrt{\beta})$. In the weak coupling expansion such terms are treated as perturbations and should be expanded. These terms are not infrared finite. And even though the final result is infrared finite it is not obvious that such terms may be considered as perturbation in the TL. Indeed, in the infinite volume limit practically all connectors become infinitely long, therefore it is not clear if the last sum is properly bounded in this limit and may be expanded perturbatively.

3. Dual representation, 't Hooft loop and monopoles

In this section we give the dual form of $SU(2)$ LGT which can be obtained from the plaquette formulation. Let x be a site dual to the cube of the original lattice and l be a link dual to the plaquette of the original lattice. Then it follows from (7), (8) and from the expression for the Jacobian that in the case of the $SU(2)$ gauge group the partition function on the dual lattice can be written in the following form

$$Z = \sum_{r_x=0, \frac{1}{2}, 1, \dots}^{\infty} \prod_x \left[(2r_x + 1) \sum_{m_i(x)=-r_x}^{r_x} \right] \prod_l \Xi_l. \quad (19)$$

The summation over r_x corresponds to the summation over all irreducible representations of the $SU(2)$ group. The sums over magnetic numbers $m_i(x)$ correspond to the calculation of $SU(2)$ traces. The index i may run from 6 to $6 + 3L$ depending on the position of the original cube, L is the linear extent of the lattice. The link integral Ξ_l is given by

$$\Xi_l = \int dV e^{\beta \text{Tr} V} V_{r_x}^{m_1 n_1} V_{r_x+n}^{\dagger m_2 n_2} \prod_{i=1}^{M(x)} \left(V_{r_i}^{k_i k_{i+1}} V_{r_i}^{\dagger p_i p_{i+1}} \right), \quad (20)$$

where V_r^{mn} is a matrix element of r -th representation. The integer number $M(x)$ depends on the position of the plaquette on the lattice and shows how many times a dual link serves as connector in the Bianchi identities.

Consider now the 't Hooft line which in 3D consists of a string S_{xy} of dual links between sites x and y on the dual lattice. With the set T of plaquettes dual to the string S_{xy} we define the action

$$S_z(U; T) = \beta \sum_{p \notin T} \text{ReTr} U_p + \beta \sum_{p \in T} \text{ReTr} z U_p \quad (21)$$

and the corresponding partition function $Z(z; S_{xy})$ with action (21). Here z is center element of $SU(N)$. By a change of variables $V_l \rightarrow z^* V_l$ for $l \in S_{xy}$

we get the following representation for the partition function $Z(z; S_{xy})$

$$Z(z; S_{xy}) = \int \prod_l dV_l \exp[\beta \sum_l \text{Re Tr} V_l] \prod_{i=1}^3 \prod_{x(i)} J(V_x^{(i)}) \\ \prod_{x(1) \neq x, y} J(V_x^{(1)}) J(z V_x^{(1)}) J(z V_y^{(1)}) , \quad (22)$$

since z cancels from all Bianchi constraints but those for the endpoints. Therefore, one can say that the Bianchi constraint is violated at these endpoints. It follows from these expressions that in the ensemble defined by the partition function (19) the disorder operator becomes particularly simple

$$\langle D_z(S_{xy}) \rangle = \frac{Z(z; S_{xy})}{Z(1; S_{xy})} = \langle \exp[2i\pi(r_x - r_y)] \rangle . \quad (23)$$

Comparing this expression with the partition function (13) and remembering that r_x is just the radial component of the auxiliary field $\alpha_k(x)$ we conclude that introducing sources like $r_x - r_y$ amounts to a simple shift in the corresponding sites for the integers m_x in (13). Therefore, we may interpret the summation over m_x as summation over monopole configurations which appear due to the periodicity of $SU(2)$ delta-function (in close analogy with $U(1)$ model). Indeed, expressed in terms of elements of algebra the Bianchi identity reads $(\sum_k \omega_k^2(x))^{1/2} = 2\pi m_x$. Configurations $m_x \neq 0$ we term monopoles of the maximal axial gauge. Investigation of these configurations and their contribution to the Wilson loops will be presented in a separate publication.

References

1. Wilson, K.G., (1974) *Phys.Rev.* **D10** p.2445.
2. Banks, T., Kogut, J., Myerson, R., (1977) *Nucl.Phys.* **B121** (1977) p.493.
3. Anishetty, R., Cheluvvaraja, S., Sharatchandra, H.S., Mathur, M., (1993) *Phys.Lett.* **B314** p.387; Halliday, I., Suranyi, P., (1995) *Phys.Lett.* **B350** p.189.
4. Borisenko, O., Faber, M., in Bogolyubov, P., Jenkovszky, L., (eds.), New trends in high-energy physics, 2000, Kiev p.221; in Lucha, W., Maung, K.M., (eds.), Quark Confinement and the Hadron Spectrum IV, 2002, World Scientific Publishing, p.269.
5. Halpern, M.B., (1979) *Phys.Rev.* **D19** p.517.
6. Batrouni, G., (1982) *Nucl.Phys.* **B208** p.467; *Nucl.Phys.* **B208** p.12.
7. Mack, G., Petkova, V., (1980) *Annal.Phys.* **125** p.117.
8. Patrascioiu, A., Seiler, E., (1995) *Phys.Rev.Lett.* **74** p.1924.
9. Borisenko, O., Voloshin, S., Faber, M., in preparation.
10. Borisenko, O., Kushnir, V., Velytsky, A., (2000) *Phys.Rev.* **D62** p.025013.
11. Müller, V.F., Rühl, W., (1981) *Annal.Phys.* **133** p.240.